

Penetrance of Arsenic Modulates Gene Expression Of Epithelial Mesenchymal Transition Marker Sox4, EPCAM and CK19 Genes Through MTHFR C677t Polymorphism Regulation in Circulating Tumour Cells Isolated from Breast Cancer Patients

Saxena AK^{1*}, Tiwari SM² and Singh P³

¹Molecular Human Genetic Laboratory, Department of Pathology/Lab Medicine, India

²Molecular Human Genetic Laboratory, Department of Biochemistry, India

³Molecular Human Genetic Laboratory, Department of Radiation Oncology, India

*Corresponding author:

Ajit Kumar Saxena,
Molecular Human Genetic Laboratory, Department
of Pathology/Lab Medicine, India

Received: 02 Mar 2024

Accepted: 22 Apr 2024

Published: 29 Apr 2024

J Short Name: COO

Copyright:

©2024 Saxena AK, This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and build upon your work non-commercially.

Citation:

Saxena AK, Penetrance of Arsenic Modulates Gene Expression Of Epithelial Mesenchymal Transition Marker Sox4, EPCAM and CK19 Genes Through MTHFR C677t Polymorphism Regulation in Circulating Tumour Cells Isolated from Breast Cancer Patients. Clin Onco. 2024; 7(10): 1-3

Keywords:

Arsenic; Epithelial mesenchymal transition marker; MTHFR C677T gene; Circulating tumor cells; Breast cancer

1. Abstract

Arsenic is a common environmental agent that modulates various kind of disease including cancer. However, the exact role of arsenic induce in breast cancer development remains unknown in circulating tumor cells (CTCs). Therefore, present study has been designed with the aims to elucidate the impact of arsenic exposure i.e 24, 48, and 72 hours on epithelial-mesenchymal transition (EMT) markers such as Sox4, EpiCAM, and CK19. RT-PCR analysis was carried out for the characterization of EMT with specific primers on agarose gel electrophoresis. DNA copy number variation (DNACNVs) was evaluated to assess genetic susceptibility and MTHFR C677T gene polymorphism was examined to determine the “risk factor” in CTCs isolated from breast cancer patients. Interestingly, findings reveal the differential gene expression and a pattern of down regulation (under expression) were observed during 24 and 48 hours of exposure, followed by up regulation at 72 hours. Genetic heterogeneity of the MTHFR CT allele was detected only at 48 hours, suggesting that arsenic increase “risk” for developing breast cancer by modulating EMT marker either alone or synergistically with MTHFR in CTCs.

2. Introduction

Arsenic, naturally occurring mutagen exist in ground water and high concentration is injurious to health (1). The continuous exposure as a source of contaminated drinking water, food preparation and irrigation of crops affecting food chain that may lead to several diseases including cancer due to chromosome aberrations and single strand DNA damage (2). There are large number of basic histopathological and cytological including immunochemistry techniques are documented in the literature whose sensitivity varying up to 60-80% and time taking too. The application of circulating tumor cells (CTCs) for early diagnosis using epithelial mesenchymal transition (EMT) is frontier in cancer medicine, whose efficiency is more than 99% in clinical diagnosis of primary solid tumors (3). Although, the etiopathology of breast cancer and interaction to environmental mutagen (As) with epithelial mesenchymal transition (EMT) during progression of disease has not been defined satisfactorily in the literature. Therefore, the present study has been designed with the aims to evaluate gene regulation of epithelial cell adhesive molecule (EpCAM) and SRY HMG-BOX transcription factor 4 (Sox4), and cytokeratin 19 (CK19), using in-vitro exposure at different time intervals. DNA copy number variation (DNACNVs) and MTHFR C677T gene polymorphism

also determined to assess the genetic susceptibility and risk factor, respectively, in circulating tumor cells isolated from breast cancer patients for early prognostic as diagnostic marker.

3. Material and Methods

Clinically diagnosed cases of breast cancer patients (n=9) and equally age matched controls were used for short term lymphocytes cultures in triplicates along with controls. Present study is approval by the Institutional Ethical Committee (IEC) of All India Institute of Medical Sciences Patna. Blood samples (0.5ml) were collected after informed consent, The cells were exposed with arsenic 0.01ug/ml (single dose) at different time intervals at 24, 48 and 72 hours, after harvesting, the circulating tumour cells '(CTCs) were isolated using Ficoll's gradient methods (4) and total cells were divided in two groups- first cells were fixed in RNazole for RNA isolation followed by gene-expression. In second group, the cells were used for DNA isolation using Promega kit (USA) and quantified by nanodrop spectrophotometer (Thermo Fisher, USA) and DNA, RNA were store at -80°C, till further study. Polymerase chain reaction (PCR) product of the EpCAM, CK-19 and SOX4 gene expression were characterized by using specific primers (forward & reverse) after confirmation by NCBI Blast. PCR reaction contains 5 µl of 5X Go Taqbuffer, 1 µl of primer set, 1.25 µl dNTP, 0.2 µl TaqDNA polymerase, 50 ng of template DNA and further volume was maintained by nuclease free water. The cycling condition was initial denaturation at 95°C for 5 minutes, denaturation at 95°C for 45s, annealing temperature based on specific target gene as documented in table-1 for 30s and extension at 72°C for 1minute, final extension at 72°C for 7 minutes. PCR products were characterized on agarose gel (1.5%) and individual bands were characterized GelDoc (Bio Red USA). DNACNVs of the EMT marker SOX4, EpCAM and CK19 genes were determined using densitometry analysis of individual band intensity and compare with controls using inbuilt Image Lab software inside GelDoc (5,6,7). Amplification-refractory mutation system (ARMS-PCR) was used for MTHFR C677T gene polymorphism using tetraplex primer forward/reverse (table-1), assay was performed using SYBR Green and Tm values were calculated to determine genetic heterogeneity between C/ T allele. The cycling conditions consist of initial denaturation at 95°C for 10 min, followed by 30 cycles of amplification steps (95°C for 10 s, 58°C for 10s, 72°C for 10s), final elongation at 72°C and followed by melt-curve analysis as detailed conditions described earlier (8). Melting curves were constructed by lowering the temperature to 65°C and later increasing the temperature by 0.2°C/s to 98°C to measuring the changes in fluorescence intensity.

4. Results

Figures-1, showing the PCR based analysis of differential expression of epithelial mesenchymal transition marker gene expression (Sox4 EpCAM and CK19), after 24, 48 and 72 hours exposure of arsenic. Similarly, bar diagram showing DNACNVs of individual

bands using densitometric analysis with controls (red). The gene expression of EpCAM and Sox4 significantly down regulated with respect to controls at 24 and 48 hours of arsenic exposure (fig.1AB). Sox4, the early transcription factor also showing a pattern of down-regulation in first 24 to 48 hours and suddenly increased at 72 hours, although the EpCAM and CK19 gene expression also up regulated at 72 hours and reach equal up to controls (fig.1C). MTHFR C677T gene polymorphism showing genetic heterogeneity at 48 hours, where, the Tm values shift from 81.50 (case) to 80.00 (GAPDH) with respect to controls.

5. Discussion and Conclusion

In the present study showing the differential expression of EMT marker gene which is modulating perhaps due to heterogeneous cell -population of breast cancer Sox4, is early transcription factor 4 (SOX4) marker belongs to gene family of Sry related high mobility group box (SOX-HMG) box region whose gene is mapped on chromosome 6p22.3 with single-exon that encodes protein 47KD comprising of 474 amino acid residues to regulates cellular differentiation and maintenance of growing progenitor cells during malignancy either alone or synergetic manner with transforming growth factor (TGF) mediating signalling during malignancy [11],The significant increase in over expression of EpCAM, CK19 and Sox4 gene and DNACNVs at 72 hours, suggesting that involving another path way to recover the damage protein (s) based on antioxidant glutathione-S-transferase gene polymerase by scavenging of free-radicals and nullify the action of toxin (arsenic) on CTCs in breast cancer patients. CK19 is associated in variety of tumour and or over expression (mRNA) was observed in primary and metastatic lesion of breast cancer patients [12-13]. EpCAM, a transmembrane cell adhesion protein and over expression is observed in both breast as well as ovarian cancer [14-15]. Earlier study of the same author is the evident that differential expression of Oct4 and Nanog and Sox2 genes fail to maintain pluripotency during organogenesis (16). Earlier study of CTCs in pancreatic tumors also reveals genetic heterogeneity of MTHFR C677T gene polymorphism (10). Interestingly, present study showing genetic heterogeneity at 48 hours due to substitution of nucleotide (point mutation) from cysteine to thymidine followed by change in amino acid from alanine to valine confirming may be due to accumulation of arsenic in first 24 hours of the cell-cycle and toxic effect on folate metabolism was observed at 48 hours in CTCs, suggesting increasing "risk factor" for breast cancer patients. These finding also supported by DNACNVs to increase genetic susceptibility either with MTHFR or EMT gene expression, hence, timely management is required through early diagnosis and becomes an essential to help for the clinicians.

6. Acknowledgement

AKS is thankfully acknowledged to the Executive Director, AIIMS Patna, for providing valuable suggestions. We also extend our thanks to the Department of Science and Technology (DST),

New Delhi for providing financial assistance and patients who participated in this study.

7. Ethics Approval

The study was approved by Institute ethical committee (IEC) of AIIMS Patna (AIIMS/Pat/IRC/2020/610).

8. Conflict of Interest

There is no conflict of interest between the authors

References

1. WHO. Guidelines for drinking-water quality, 4th edition incorporating the first addendum. Geneva, World Health Organization. 2017; 315–318.
2. Emma G, Amadeu C, Jessica P, Enrico S, et al. In-vitro DNA damage by arsenic compounds in a human lymphoblastoid cell line (TK6) assessed by the alkaline Comet assay *Mutagenesis*. 2004; 19(2): 129–135.
3. Ankeny JS, Court CM, Hou S, Li Q, Song M, Wu, D. Tomlinson, JS. Circulating tumour cells as a biomarker for diagnosis and staging in pancreatic cancer. *British journal of cancer*. 2016; 114(12), 1367-1375.
4. Saxena Ajit K, Shalini, Tiwari M, Singh P and Singh V. Comprehensive mutational spectra of epithelial mesenchymal transition markers-SOX4, EpCAM, CK19 and their impact of methylenetetrahydrofolate reductase C677T gene polymorphism associated risk factor modified by BRCA1/2 and TGF- β R1 genes in pre/postmenopausal cases of breast cancer, *International Journal of Cancer Medicine*. 2024; 7(1); 9-29.
5. Zhang J, Liang Q, Lei Y, Yao M, Li L, Gao X, et al. SOX4 induces epithelial–mesenchymal transition and contributes to breast cancer progression. *Cancer Res*. 2012; 72: 4597–608.
6. Alowaidi F, Hashimi SM, Alqurashi N, Alhulais R, Ivanovski S, Bellette B, et al Assessing stemness and proliferation properties of the newly established colon cancer stem cell line, CSC480 and novel approaches to identify dormant cancer cells. *Oncol Rep*. 2018; 39:2881–91.
7. Balducci E, Azzarello G, Valori L, Toffolatti L, Bolgan L, Valenti MT. A new nested primer pair improves the specificity of CK-19 mRNA detection by RT-PCR in occult breast cancer cells. *Int J Biol Markers*. SAGE Publications Sage UK: London, England; 2005; 20:28–33.
8. Saxena A K, Gupta RK. ARMS-PCR based SNP analysis of MTHFR C677T allele using Syber green in pancreatic tumour. *British Journal of Medicine and Medical Research*. 2016; 11(12): 1-6.
9. Song GD, Sun Y, Shen H, Li W. SOX4 over expression is a novel biomarker of malignant status and poor prognosis in breast cancer patients. *Tumour Biology*. 2015; 36(6): 4167–4173.
10. Saxena A K. Disconcordance Between stem cell Oct4, Sox2, Sox4 and epithelial mesenchymal transition CK 19, EpCAM markers in circulating tumour cells-MTHFR C677T gene variant increase risk factor in pancreatic tumours. *Journal of Cancer Medicine & Metastasis*. 2023; 6(1): 1-9.
11. Moreno CS. SOX4: The unappreciated oncogene. In *Seminars in cancer biology*. 2020; 67: 57-64.
12. Keyvani S, Karimi N, Orafa Z, et al. Assessment of cytokeratin-19 gene expression in peripheral blood of breast cancer patients and breast cancer cell lines. *Biomarkers in Cancer*. 2016; 8: BIC-S38229.
13. Orafa Z, Karimi N, Keyvani S, et al. Quantitative CK19 biomarker detection in breast cancer cell lines. *Journal of Medicine and Life*. 2022; 15(2): 188-195.
14. Litvinov SV, Bakker HA, Gourevitch MM, et al. Evidence for a role of the epithelial glycoprotein 40 (Ep-CAM) in epithelial cell-cell adhesion. *Cell Adhesion and Communication*. 1994; 2(5): 417-428.
15. Schmidt M, Hasenclever D, Schaeffer M, et al. Prognostic effect of epithelial cell adhesion molecule overexpression in untreated node-negative breast cancer. *Clinical Cancer Research*. 2008; 14(18): 5849-5855.
16. Went P, Vasei M, Bubendorf L, et al. Frequent high-level expression of the immunotherapeutic target Ep-CAM in colon, stomach, prostate and lung cancers. *British Journal of Cancer*. 2006; 94(1): 128-135.
17. Saxena, A K. Shalini, Singh V, Singh P. Differential Expression of Vimentin Gene Modulated by MTHFR Gene Polymorphism in Circulating Tumor Cells Isolated from Breast Cancer Patients. *Genetics and Molecular*. 2024; 23 (1): 1-10.